

The History of Asteroid Family Identification

Zoran Knežević

Serbian Academy of Sciences and Arts
Kneza Mihaila 35, 11000 Belgrade, Serbia
email: zoran.knezevic@sanu.ac.rs

Abstract. In this paper the early history of search for asteroid groupings is briefly reviewed. Starting from the first attempts by Kirkwood, who identified a number of asteroid pairs and triples with adjacent orbits, via the similar contributions of Tisserand and Mascart, we arrive to Hirayama and his discovery of asteroid families, marking the beginning of modern asteroid science.

Keywords. history of astronomy, asteroid pairs, asteroid families

Asteroid families are discovered a century ago by K. Hirayama. The seminal paper of his: “Groups of asteroids probably of common origin” (Hirayama 1918), as well as a series of subsequent papers devoted to asteroid families (Hirayama 1919, 1920, 1922, 1927, 1933), received a lot of attention in the past, and have been thoroughly described in a recent review (Knežević 2016). In the present paper, we shall therefore pay more attention to several almost forgotten attempts to find asteroid groupings that precede the Hirayama’s discovery.

The first attempts to find groups of asteroids with similar orbits date back to 1877, when D. Kirkwood, in an article entitled “The asteroids between Mars and Jupiter” (Kirkwood 1877), presented ‘evidence of a similarity more than accidental between adjacent orbits of the asteroidal group’. Among only 172 asteroids known at the time, he found 4 pairs (Fortuna and Eurynome, Fides and Maia, Clotho and Juno, Sirona and Ceres) with striking similarity in the magnitude, form and position of orbits. For an additional pair (Urda and Gerda) he states that the discoverer of these bodies, Dr. Peters, in the *American Journal of Science* for February 1877, calls attention to coincidences between the elements of their orbits. Kirkwood does not venture (yet) to speculate on the origin of such pairs, stating simply that they must be of common origin.

In 1888 Kirkwood publishes a booklet (Kirkwood 1888) devoted entirely to asteroids, in which he again mentions ‘adjacent orbits’, drawing, however, a special attention to the pair Hilda and Ismene, for which he finds distances, periods, inclinations and eccentricities nearly identical, but longitudes of perihelia different by almost exactly 180° . Wondering whether this indicates that the two objects detached at about the same time from the opposite sides of the solar nebula, he suggests that in general the asteroid pairs with similar orbits originate from single asteroids, which, like comets, ‘may have been separated by the sun’s unequal attraction on their parts’.

Finally, in a couple of papers published in astronomical journals (Kirkwood 1890, 1891) he again returns to the topic of similarity of asteroid orbits, proposing in the former paper the three most significant pairs (including a new one – Vera and Semele) and again discussing possible formation by separation/dismemberment; in the latter paper, however, starting from the catalog with about 300 asteroids, he reports on finding of 10 groups (4 triples and 6 pairs) of bodies with similar orbits. Assuming certain characteristics of the solar nebula, he concludes that the disturbing influence of Jupiter could

have been sufficient not only to detach the masses of asteroids from the central body, but also to subdivide by unequal attraction on the different parts the newly-formed nebulous planets, until the fragments finally resulted in the existing asteroids.

In his article “La question des petites planètes”, F. Tisserand (1891) reports ‘sans explication’ on the three pairs of asteroids with similar orbital elements: 106 Dione and 245 Vera (forming with 86 Semele one of the triples proposed by Kirkwood), 218 Bianca and 246 Asporina, and 84 Clio and 249 Ilse (again a pair which makes part of the Kirkwood’s triple with 115 Thyra).

Finally, J. Mascart published two studies (Mascart 1899, 1902) in which he considers pairs of asteroids with similar orbits. In the former paper, he first computes Tisserand invariants for a total of 417 asteroids, identifying subsequently 20 pairs of asteroids with very close values of the invariants and with more or less similar orbital elements. In the latter paper he then analyses in detail the probabilities of these coincidences.

As pointed out by Hirayama (1922), all these attempts to find significant groupings of asteroids failed. Hirayama first rightly cautions against having only two or three asteroids with similar orbits, because this gives rise to a considerable probability that they may be accidentally coincident. Then, explaining the failure to find more asteroids with similar orbits, he attributes it to the ‘use of actual orbits ... for comparison, whereas these orbits are varied remarkably by the action of the planets’, and correctly states that such attempts could be successful only if the separation had occurred very recently. To identify families originated at remoter age, Hirayama asserts that some kind of invariable elements must be used to identify families. Indeed, in the same paper he formally introduces the *proper elements* as such invariable parameters and computes them by means of the Lagrangian linear theory of secular perturbations, stating: ‘... if there may be noticed a group of the asteroids with the elements approximately common to all, and if at the same time, their number is sufficiently large to be insured from the effect of chance, then we can conclude that they have probably originated from the breaking up of a single asteroid’. His success in identifying asteroid families with tens of members marks the end of the pioneer era of the search for asteroid groupings, and the beginning of the modern asteroid science.

References

- Hirayama, K. 1918, *Astron. J.*, 31, 185
 Hirayama, K. 1919, *Proc. Phys. Mat. Soc. Japan*, III, 1, 52
 Hirayama, K. 1920, *Proc. Phys. Mat. Soc. Japan*, III, 2, 236
 Hirayama, K. 1922, *Japanese J. Astron. Geo.*, 1, 55
 Hirayama, K. 1927, *Japanese J. Astron. Geo.*, 5, 137
 Hirayama, K. 1933, *Proc. Imp. Acad. Japan*, 9, 482
 Kirkwood, D. 1877, *The Annual Review of the Board of Regents of the Smithsonian Institution for 1876* (Washington: Gov. Print. Office) p. 358
 Kirkwood, D. 1888, *The Asteroids or Minor Planets Between Mars and Jupiter* (Philadelphia: J.B. Lippincot Comp.)
 Kirkwood, D. 1890, *Publ. Astron. Soc. Pacific*, 2, 48
 Kirkwood, D. 1891, *Publ. Astron. Soc. Pacific*, 3, 95
 Knežević, Z 2016, in: S.R. Chesley, A. Morbidelli, R. Jedicke & D. Farnocchia, (eds.) *Proceedings IAU Symposium 318: Asteroids: New Observations, New Models* (Cambridge: Cambridge Univ. Press), p. 16
 Mascart, J. 1899, *Bull. Astron.*, 16, 369
 Mascart, J. 1902, *Annales Obs. Paris*, XXIII, F
 Tisserand, F. 1891, *Annuaire Pour L’An 1891 publie par Bureau des Longitudes* (Paris: Gauthier-Villars et files) p. B.18